HYDROMIN BOTON 20/0 Amino Acid Complex

GUARNANTEED ANALYSIS

Boron (B) _____2.0%

Derived from: Sodium Borate. Complexing Agent from Hydrolyzed Protein

NET CONTENTS 5 GALLONS 18.93 LITERS 9.5 LBS. PER GAL @ 68° F 1062 GRAMS PER LITER @ 20° C

KEEP OUT OF REACH OF CHILDREN

WARRANTY: Western Nutrients Corporation makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on the Western Nutrients Corporation Label attached to the product container

Nutraplex is a Registered product of Western Nutrients Corporation

PRODUCT INFORMATION CROPS

HYDROMIN BORON can be applied to most vegetable crops, row crops, deciduous fruit and nut trees, citrus, avocados, grapes, melons, ornamentals, turf, pasture, range grasses, and most other crops.

HYDROMIN BORON liquid foliar fertilizer was formulated to provide a fast method to boost plants during critical or high growth periods, damage due to weather and for correction of nutrient deficiencies.

HYDROMIN BORON promotes seed and root development as well as a strong cell structure when used as a supplement to a soil fertilizer program. Micro-Nutrients are needed for healthy seed development, root structure and cell structure, especially in soils where the nutrients have been depleted in sufficient quantities to cause reduced yields or stunted plants. HYDROMIN BORON insures against these nutrient deficiencies providing the essential nutrients to stimulate early root development and speed up crop growth.

HYDROMIN BORON is a partially chelated liquid micronutrient for foliar and soil application to agricultural crops. HYDROMIN BORON is plant food and non-phytotoxic when used as directed. It is absorbed through the leafy tissue and root system of the plant. HYDROMIN BORON is compatible with most insecticides, fungicides, herbicides, liquid fertilizer, and other foliar nutrients. It is used on most field and row crops, trees, vines, turf, and ornamentals.

HYDROMIN BORON can be applied by conventional ground rig, concentrate sprayer, and by air. For additional micro nutrients add 1-3 quarts (1-2 liters) per acre of NUTRAPLEX® products with HYDROMIN BORON.

MANUFACTURED BY - WESTERN NUTRIENTS CORPORATION

245 Industrial Street, Bakersfield California 93307 • (661) 327-9604
(661) 327-1740 Fax • (800) 542-6664 Ca. Only
E-mail: info@westernnutrientscorp.com
Website: http://www.westernnutrientscorp.com

SUGGESTED USES

GENERAL USE- 1-3 quarts (1-3 liters) per acre.

May be applied by air, concentrate sprayer, or conventional ground rig. Maximum concentration 1 quart (1 liter) HYDROMIN in 5 gallons of solution. HYDROMIN should be used in regular crop protection spray programs to insure plants receive all the essential plant foods

ALFALFA - 1-2 quarts (1-2 liters) per acre after each cutting for hay. On seed alfalfa, 4-6 applications.

ARTICHOKES - 1-2 quarts (1-2 liters) per acre at bud stage. Repeat every 2-3 weeks.

AVOCADOS, CITRUS - 1-2 quarts (1-2 liters) per acre pre-bloom.

BLACKBERRIES, BOYSENBERRIES, RASPBERRIES, STRAWBERRIES ands other small fruits - 1 quart (1 liter) per acre pre-

BROCCOLI, BRUSSEL SPROUTS - 1-2 quarts (1-2 liters) per acre 4-6 applications.

CARROTS, ONIONS, PEPPERS, GARLIC, POTATOES, SPINACH - 1-2 quarts (1-2 liters) per acre as often as needed.

CAULIFLOWER, CABBAGE - 1-2 quarts (1-2 liters) per acre, 4-6 applications.

CORN - 2 quarts (2 liters) per acre. Apply with crop protection sprays at 7-10 day intervals. Begin 3 weeks after emergence.

COTTON - 1 quart (1 liter) per acre prior to bloom. Make 4-6 applications.

CUCUMBERS - 1-2 quarts (1-2 liters) per acre pre-bloom. Make 4-6 applications. Apply with crop protection sprays

DECIDUOUS FRUIT and NUT TREES - 1-3 quarts (1-3 liters) per acre pre-bloom and petal fall.

GRAPES - 1-2 quarts (1-2 liters) per acre prior to bloom. Make 3 applications.

LETTUCE, CELERY - 1 quart (1 liter) per acre as often as needed.

MELONS, PEAS, and BEANS, SQUASH - 1-2 quarts (1-2 liters) per acre, pre-bloom. Make 4-6 applications.

SOYBEANS - 2-3 guarts (2-3 liters) per acre prior to bloom and after pod formation.

SUGAR BEETS - 2 quarts (2 liters) per acre. Make several applications

TOBACCO - 1-2 quarts (1-2 liters) per acre. Apply with crop protection sprays.

TOMATOES, EGGPLANT - 1 quart (1 liter) per acre at first bloom. Repeat before and during fruiting period.

WHEAT, BARLEY, MILO, RICE and other GRAIN CROPS - 1-2 quarts (1-2 liters) per acre seeding and pre-bloom.

LAWN and TURF - 1-2 quarts (1-2 liters) to 100 gallons (378 liters) of water per 5000 square feet.

NURSERY and GREENHOUSE - 1 quart (1 liter) per 100 gallons (378 liters) of water.

TRANSPLANTING SOLUTIONS - 6 ounces per 10 gallons (37.8 liters) solution.



This product can expose you to chemicals including Cobalt Sulfate, which is known to the State of California to cause birth defects or other reproductive harm.

For more information go to www.P65Warnings.ca.gov